

Charge densities above pulsar polar caps

Axel Jessner¹, Harald Lesch² and Thomas Kunzl^{2,3}

ABSTRACT

The ability of the neutron star surface to supply all or only part of the charges filling the pulsar magnetosphere is crucial for the physics prevailing within it, with direct consequences for the possible formation of pair creation regions. We evaluate the conditions for e^- emission from pulsar surfaces for a simple Goldreich-Julian geometry taking both thermal and field emission processes into account. Using recently published estimates for the equation of state at the neutron star's surface, we show, that for a large range of T_{surf} , B and P , the liberated charges will fully screen the accelerating B-parallel electric field $E_{\parallel}$. For surface temperatures $T_{\text{surf}} < 2 \cdot 10^5 \text{K}$ a balance between field emission of electrons and shielding of the field will occur. Even in the overidealised case of $T_{\text{surf}} = 0$ one can expect a prodigious supply of electrons which will weaken the accelerating $E_{\parallel}$. We calculated the motion of electrons along selected polar field lines numerically for the low temperature, field emission scenario yielding their Lorentz factors as well as the produced radiation densities. Inverse Compton and later curvature losses are seen to balance the acceleration by the residual electric fields. We found that the conditions for magnetic pair production are not met anywhere along the field lines up to a height of 1500 pulsar radii. We did not *a priori* assume an "inner gap", and our calculations did not indicate the formation of one under realistic physical conditions without the introduction of further assumptions.

Subject headings: Plasmas - Radiation mechanism - Pulsars

¹Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany

²Universitäts-Sternwarte München, Scheinerstr.1, D-81679 München, Germany

³Max-Planck-Institut für extraterrestrische Physik, Giessenbachstr., D-85740 Garching, Germany

1. Introduction

A proper assessment of the conditions very close to the surface of a neutron star is a vital ingredient for any model that attempts to explain the enigmatic origin of the observed electromagnetic radiation from pulsars.

Numerous studies of the polar cap acceleration have been made within a particular framework: A steady acceleration of electrons within a space charge limited particle flow originating at the surface of the neutron star will produce particles emitting magnetically convertible curvature γ -rays (e.g. Arons & Scharlemann 1979). These high energy photons are converted into e^+e^- -pairs with high enough efficiency (Daugherty & Harding 1982) so that in the end, they and their daughter products will exceed the initial (Goldreich-Julian density) by a factor of 10^4 (e.g. Zhang & Harding 2000). An important *a priori* assumption is inherent in all these calculations: Above a critical height h_{gap} a region of high conductivity will quench any electric field $E_{\parallel}$ parallel to the guiding B-field lines, so that the full potential across the polar cap will be active between the outer surface of the neutron star and the inner surface of the high conductivity region.

Our approach differs in that respect from other workers, we do not make that initial assumption, but try to calculate the amount of charges produced by the surface, how they affect and are affected by $E_{\parallel}$, their energy losses and the possibility of pair production. Should one find suitable conditions for such a pair production region to exist, the methods quoted above will be the prime choice for its description.

The conclusion that electron-positron pairs with high densities exist close to the neutron star has recently been questioned by several authors. Kunzl et al. (1998) considered this widespread assumption of dense pair plasmas for the problem of radio wave propagation within the neutron star magnetosphere. On hand of the most recent emission height estimates (50 - 100 R_{NS}) for the radio emission (Kijak & Gil (1997); Kramer et al. (1997, 1998)) they found that the existence of low frequency radio emission for example from the Crab pulsar is incompatible with the scenario of extended dense pair production. These findings were fully confirmed by the theoretical investigation of Melrose & Gedalin (1999), who treated the plasma emission mechanisms in detail and came to the conclusion that only low energy particles with

Goldreich-Julian density could exist in the radio emission regions in order to allow for escaping radio waves (see also Melrose (2000) and references therein). Furthermore recent simulations of the pair cascade by Arendt & Eilek (2000) for strong and weak magnetic fields shed some light on the difficulties for an efficient production of secondary pairs by high energetic primary particles via γ -ray curvature photons. Although their findings need some more investigation they could show that the field strength at the surface has the most important effect on the cascade. For strong fields $B \sim 10^9$ T they find almost no secondary photons and the cascade terminates in one generation. The weak fields ($B \sim 10^7$ T) had either no cascades or quite weak ones. Only for moderate fields ($B \sim 10^8$ T) they observed some secondary particle generations. Arendt & Eilek concluded that it is not yet clear whether pulsars can really produce a large number of pairs at the surface gaps.

This tempted us to question the canonical *a priori* assumption of the existence of an inner pair creating gap and investigate its possibility given a reasonable estimate of the surface characteristics and the fields involved.

In the literature there are three fundamentally different ideas for pair production in the inner magnetosphere: A "starved" magnetosphere where the neutron star surface provides only an insufficient supply of charged particles (e.g. Ruderman and Sutherland (1975)), a magnetosphere where the Goldreich-Julian density is reached at the surface but due to curvature effects of the field lines deviations occur close to the surface (e.g. Arons & Scharlemann (1979), Barnard & Arons (1982)) and Shibata's model which predicts electric fields caused by *overdense* regions, that means in the opposite sense than in the Arons model (Shibata, 1997).

The latter two ideas suffer from serious weaknesses since it turned out that Arons type models do not work for slow pulsars (Beskin (2000) gives an upper limit of 0.1 to 0.3 s for the period). The Shibata model in turn neglects the back reaction of an overdense region which reduces the particle flow from inside. Therefore we will restrict our discussion to models of Ruderman-Sutherland type and fix the parameter range in which this kind of models is reasonable.

For the scope of this article we consider a strong magnetic dipole anchored in a heavy ($1.4M_{\text{sun}}$) sphere with a radius $r = 10$ km that rotates twice per second ($\Omega = 2\pi \cdot 2s^{-1}$). Let the B-field have a strength of

$|B_0| = 10^8$ T at the surface and for simplicity's sake let the magnetic and the rotational axes be aligned¹.

$$B(r, \theta) = B_0 \cdot \left(\frac{r_{\text{NS}}}{r}\right)^{-3} \cdot (\cos \theta, \frac{1}{2} \sin \theta, 0) \quad (1)$$

Such objects have been discussed as idealised cases of radio pulsars by Goldreich & Julian (1969) and many authors since then. Of course one has to realise that the above mentioned idealisations limit the applicability of such simple models for the interpretation of observational facts from the growing number of pulsars that is being discovered. We will however try to extend the scope of these models by constraining the number of charged particles available and analysing the fate of individual charged particles (limiting ourselves to electrons and positrons) in the polar regions of such heavy rotating dipoles under the constraints given by some of the more recent observations. Although a pulsar magnetosphere is characterized by extreme physical conditions in terms of strong electric and magnetic fields the fundamental processes that can supply charges are reasonably well known from terrestrial experiments and technical applications and can be extrapolated to the surface conditions of a neutron star. Such an extrapolation provides constraints on the initial charge density which is vital for any model of cascading particle creation and the radiation processes.

2. Accelerating fields and charge densities

Muslimov & Tsygan (1990, 1992) showed how the electric and magnetic fields of a rotating neutron star are distorted by relativistic effects. A comprehensive treatment of the rotating dipole in the frame of general relativity was given by Muslimov & Harding (1997). For our purposes of *demonstrating* the major physical properties near the polar cap it suffices however to stick to a classical description (including special relativity of course).

Our rotating magnetic dipole acts as an unipolar generator and induces an electric field as the consequence of its surface charge distribution

$$E(r, \theta) = -B_0 \Omega r_{\text{NS}} \left(\frac{r}{r_{\text{NS}}}\right)^{-4} \cdot (3 \cos^2 \theta - 1, 2 \cos \theta \sin \theta, 0). \quad (2)$$

¹We use spherical coordinates (r, θ, ϕ) centered in the neutron star with θ being the directional angle referred to the pole which coincides with the rotation axis.

The component parallel to the magnetic field lines $E_{\parallel} = \frac{\vec{E} \cdot \vec{B}}{|\vec{B}|}$ is given by

$$E_{\parallel}(r, \theta) = -4B_0 \Omega r_{\text{NS}} \left(\frac{r}{r_{\text{NS}}}\right)^{-4} \frac{\cos^3 \theta}{\sqrt{3 \cos^2 \theta + 1}} \quad (3)$$

and reaches $2.5 \cdot 10^{13}$ V/m at the surface near the magnetic poles (Goldreich & Julian 1969; Holloway 1973). In vacuum all appropriately charged particles will be quickly accelerated to very high energies, but because of the strong magnetic field they are constrained to move along the magnetic field lines given by the parametric equation

$$r(\theta, \theta_0) = r_{\text{NS}} \cdot \frac{\sin^2 \theta}{\sin^2 \theta_0}. \quad (4)$$

Here θ_0 is meant to be the colateral angle of the field line at the surface of the sphere.

Goldreich & Julian (1969) already calculated the electron density which shields the electric field and yields the only stable static solution for a pulsar magnetosphere (excluding centrifugal and gravitational forces).

$$n_{\text{GJ}}(r, \theta) = \frac{\varepsilon_0 \Omega B_0}{e} \left(\frac{r}{r_{\text{NS}}}\right)^{-3} \cdot (3 \cos^2 \theta - 1) \quad (5)$$

with e being the electron charge.

In our example this amounts to $n_{\text{GJ}0} = \frac{2\varepsilon_0 \Omega B_0}{e} = 1.4 \cdot 10^{17} \text{m}^{-3}$ at the poles on the surface. Deutsch (1955) was the first who showed that the high induced fields make it unlikely that stellar rotators with strong magnetic fields will rotate in a (charge-free) vacuum. Several well studied processes enable i.e. electrons to escape from a metallic surface. If the neutron star surface can thus liberate sufficient charges to keep the Goldreich-Julian current flowing, the emission will be space-charge limited like an electronic valve (diode). Electrons will be accelerated away² from the polar regions if $\vec{\Omega} \cdot \vec{B} > 0$ ³.

²The observed high brightness temperatures of the pulsar radio emission suggest the operation of beaming mechanisms for the enhancement of the radio intensity. We'll therefore focus our attention on outflowing particles where intensity enhancement by forward beaming would be natural.

³In some models in the literature we need *anti-alignment* of $\vec{\Omega}$ and $\vec{B}$. In that case the Goldreich-Julian charge density above the polar cap is positive. We do not discuss this case here but since the binding energies of ions are assumed to be comparable to those of electrons the qualitative argumentation remains unchanged.

Usov & Melrose (1995) have reminded us of the fact that the relevant fundamental surface emission processes are field emission and thermal emission (perhaps enhanced by the Schottky effect (Schottky 1914)). Thermal surface emission of electrons was discovered by T.A. Edison (1884) and has been extensively studied since the beginning of our century (Dushman 1930). Field emission from metal surfaces is also known for a long time (Fowler & Nordheim 1928) and used in many technical applications. The effectiveness of both emission processes is controlled by E_W , the work function or binding energy, the energy required to liberate an electron from the crystal lattice at the surface boundary. Values for E_W and the Fermi-energy E_F for the pulsar surface can only be extrapolated from terrestrial values as the high densities of $\rho_{Fe} \approx 10^6 \text{ kg m}^{-3}$ and magnetic fields $B \approx 10^8 \text{ T}$ cannot be reached in laboratory experiments. We will therefore first try to establish a *reasonable range of values* for E_W in order to investigate the effectiveness of surface emission.

2.1. Constraints for the range of the work function

Using a Thomas-Fermi-Dirac-Weizsäcker approximation Abrahams & Shapiro (1992) calculated the cold ($T = 0$) surface density ρ_{Fe} to be about $2.9 \cdot 10^6 \text{ kg m}^{-3}$. That result enables us to give an improved estimate of the Fermi energy at the surface:

$$E_F(\rho_{Fe}) = \frac{2 \cdot \pi^4 \hbar^4 c^2}{e^2 B^2 m_e} \cdot \left(\frac{\rho_{Fe} \cdot (28 - 2)}{56 \cdot m_p} \right)^2 \quad (6)$$

which amounts to $E_F = 4.17 \cdot 10^2 \text{ eV}$ (as usual $\hbar = h/2\pi$). In our subsequent calculations we take that as the most likely value of E_F . In the case of an unknown work function E_W , one usually approximates $E_W = E_F$. So far calculations of the actual electron work function have not been made and for want of a better alternative we will use the Fermi energy as the best available measure of the work function. Abrahams & Shapiro (1992) give the ionisation energy for iron as 120 eV and we will consider the range from ionisation energy to cyclotron energy (0.12 - 12 keV) to be also a reasonable constraint for E_W .

2.2. Field (cold cathode) emission

By calculating the transmission coefficient of electron wave functions through a potential wall

$U(x) = e^{-1} E_W - E_0 \cdot x$ for $x > 0$ and $U(x) = 0$ for $x < 0$ (E_0 being the electric field) Fowler & Nordheim (1928) provided a description of electron currents from metal surfaces in good agreement with the experimental results and useful in many technical applications since then. Using their expression of the cold cathode current density and assuming a relativistic flow of electrons we find for the particle density

$$n_{\text{field}}(E_0) = \frac{e^2}{2\pi\hbar c} \frac{\sqrt{\frac{E_W}{E_F}} \cdot E_0^2}{(E_W + E_F)} e^{-\frac{8\pi \cdot \sqrt{2 \cdot m_e}}{3\hbar e} \cdot \frac{E_W^{3/2}}{E_0}} \quad (7)$$

In Fig. 1 we plot the ratio of the field emission charge density to the Goldreich-Julian density for a standard pulsar. The authors like to emphasize the point that this density is independent of the somewhat uncertain surface temperature and depends only on E_0 .




Fig. 1.— Cold cathode emission n/n_{GJ} versus work function E_W for a standard pulsar (see text).

Field emission can supply the Goldreich-Julian density for a cold neutron star surface for any $E_W < 2 \text{ keV}$! Fig. 2 however shows, that for the extreme case of a 100 times weaker field (i.e. a slowly rotating weak pulsar) the maximum binding energy drops down as far as 50 eV.

Fig. 3 shows how for an assumed work function of i.e. 500 eV, the cathode efficiency increases dramatically with the electric field. We may therefore expect




Fig. 2.— The critical work function for the Goldreich-Julian density versus the surface field E_0 .

field emission to play a significant role in the supply of charges for the average pulsar magnetosphere.

2.3. The thermal electron emission

In contrast to the quantum mechanical effect of field emission, the thermal surface emission of electrons from a conductor is a purely classical effect: For a given particle distribution function, be it Maxwellian or Fermi–Dirac, there will be a fraction of particles with a kinetic energy that exceeds that of the potential barrier characterised by the binding energy. That number of electrons liberated from a hot metallic surface is given by Dushman’s equation (Dushman 1930). In the presence of strong external electric fields, the number of electrons is increased over that given by Dushman’s equation because of the Schottky effect (Schottky 1914) which lowers the effective potential barrier for the particles. The charge density is thus described by the combined Dushman-Schottky-Equation

$$n_{\text{DS}}(T, E_0) = \frac{A_0}{e \cdot c} \cdot T^2 \cdot e^{-\frac{E_W}{kT} + \frac{e}{kT} \cdot \sqrt{\frac{e \cdot E_0}{4\pi\epsilon_0}}} \quad (8)$$

(k being the Boltzmann constant) with $A_0 = \frac{4\pi \cdot m_e \cdot e \cdot k^2}{h^3} = 120.2 \text{ Acm}^{-2}$ which depends only on E_W . In Fig. 4 we show the contributions to the current density




Fig. 3.— The cold cathode efficiency n/n_{GJ} at $E_W = 417 \text{ eV}$ as a function of the surface field E_0 .

according from field emission and from thermal emission with and without the Schottky effect for a range of temperatures likely for a standard pulsar with an assumed $E_W = 417 \text{ eV}$.

As has been pointed out before, for $E_0 = 2.5 \cdot 10^{13} \text{ V/m}$ field emission alone can supply all that is required, but for temperatures above 10^5 K , thermal emission, which is only weakly dependent on the effective field becomes even more efficient. From $2 \cdot 10^5 \text{ K}$ onwards, thermal electrons, emitted from the surface even without a large electric field can supply the GJ-density. Fig. 5 brings everything into perspective by outlining the extremes of the parameter space.

We show the effective ratio of total charge emission for work functions of $E_W = 2065 \text{ eV}$ and $E_W = 417 \text{ eV}$ as well as for a field of $\frac{E_0}{100}$. Only for extremely high $E_W \gg 1200 \text{ eV}$ or very low fields $E < 10^{12} \text{ V/m}$ and low surface temperatures $T < 2 \cdot 10^5 \text{ K}$ the electron supply from the pulsar surface will be insufficient to fill the magnetosphere with charges to shield the induced electric field. It is however doubtful that objects with such low temperatures can be found as internal stresses caused by the rotational braking of the neutron star set a lower limit of a few 10^5 K for the surface temperature (Tsuruta 1998 and references therein). But normally the potential supply of charges exceeds the requirements of the G-J-density by a wide margin. Thus any accelerating electric field at the surface of the neutron star will be quenched within an extended corotating charged lower region of the magnetosphere. Therefore we do not expect any




Fig. 4.— The surface efficiency n/n_{GJ} at $E_W = 417\text{eV}$ for a standard pulsar as a function of the surface temperature. Thin solid line: thermal emission, dots: thermal emission with Schottky effect, Thick solid: cold cathode emission.

significant acceleration in the inner magnetosphere which, of course, implies the absence of highly relativistic particles and no pair production at all under these circumstances.

3. Neutron stars dominated by field emission

We have shown in the last section, that high binding energies or low surface temperatures are the only conditions compatible with significant surface electric field strengths that can accelerate particles. In these cases, an equilibrium field of $E_{\text{eq}} = E_0 \cdot (1 - \frac{n_e}{n_{GJ}})$ will remain when n_e electrons flow off the surface. Both the electric field $E_{\parallel}$ and the shielding density are proportional to the product $B \cdot \Omega$. We introduce a scale factor $\xi = \frac{B \cdot \Omega}{(B \cdot \Omega)_0}$ to account for the possible empirical variation of $B \cdot \Omega$ and by numerically solving the balance equation

$$\begin{aligned} \xi n &= n_{\text{field}}(\xi E(1 - \frac{n}{\xi n_{GJ}}), E_W) + \\ &+ n_{\text{DS}}(\xi E(1 - \frac{n}{\xi n_{GJ}}), E_W, T) \end{aligned} \quad (9)$$




Fig. 5.— The surface efficiency n/n_{GJ} for different E_W and as a function of the surface temperature. long dash: total efficiency for $E_W = 417\text{eV}$, solid: total efficiency for $E_W = 417\text{eV}$ and $0.1E_0$, short dash: total efficiency for $E_W = 2065\text{eV}$, dots: total efficiency for $E_W = 417\text{eV}$ and a weak field pulsar with $0.01E_0$ and $0.01n_{GJ}$.

for $0.02 < \xi < 5$ we obtain the equilibrium densities for any given temperature T and work function E_W covering a wide range of neutron stars with different B and Ω . Fig. 6 shows the variation of the residual electric field $\xi E(1 - \frac{n}{\xi n_{GJ}})$ as a function of ξ for $E_W = 417\text{eV}$ and $E_W = 1200\text{eV}$. A temperature of $1.85 \cdot 10^5\text{K}$, just below the onset of thermal emission for $E_W = 417\text{eV}$, was chosen for the calculation. The residual field grows with ξ until it saturates at $2.1 \cdot 10^{12}\text{V/m}$ for $\xi = 1$. A higher E_W causes earlier saturation at a higher level. Between 300 eV and 2000 eV the residual field saturation level E_{lim} depends nearly linearly on E_W and can be approximated as $E_{\text{lim}} \approx 1.3 \cdot 10^{10} \cdot (E_W - 250\text{eV})\text{V/m}$. For our standard E_W , the density n deviates by only 10% from the Goldreich-Julian value, but higher E_W show lower initial densities with the expected increase in charge density with increasing electric fields.

3.1. The equilibrium energy of electrons in the magnetosphere

Any residual field will quickly accelerate the electrons to relativistic energies, balanced by the braking due to various energy loss processes. Because of the high magnetic field strength, the motion of charges will be along the field lines. With the help of the parametrisation (4) we can describe a particle tra-




Fig. 6.— The residual electric surface field $\xi E(1 - \frac{n}{n_{GJ}})$ at $T = 1.85 \cdot 10^5 \text{K}$ as a function of ξ for $E_W = 417 \text{eV}$ (solid line) and $E_W = 2065 \text{eV}$ (dots).

jectory using only its co-lateral angle at the pulsar surface (starting angle θ_s) and the co-lateral angle θ on the trajectory. At each point of such a trajectory, the energy gain of a charge by the parallel component of the residual electric field

$$\frac{d\gamma}{dt}|_E = - \left(\frac{e}{m_e c} \right) \left(1 - \frac{n_e}{n_{GJ}} \right) E_{\parallel}(\theta, \theta_0) \quad (10)$$

will be balanced by a number of energy loss mechanisms. Curvature radiation is an obvious loss mechanism for relativistic charged particles forced to follow a curved trajectory. The energy loss of a relativistic particle due to curvature radiation is given by (e.g. Zheleznyakov 1996, p.231)

$$\frac{d\gamma}{dt}|_{CR} = \frac{2}{3} \gamma^4 \frac{e^2}{4\pi\epsilon_0 m_e c} \frac{1}{R_C(\theta, \theta_0)^2} \quad (11)$$

Here we use the curvature radius of a dipolar field line in its parameterised form (Lesch et al. 1998)

$$R_c(\theta, \theta_0) = \frac{r_{NS}}{3} \frac{\sin \theta}{\sin^2 \theta_0} \frac{(1 + 3 \cos^2 \theta)^{3/2}}{1 + \cos^2 \theta} \quad (12)$$

The curvature radii in the polar regions of a neutron star with a dipolar magnetic field are of the order of

10^6m . To balance the huge acceleration of an unshielded field at the surface $\frac{d\gamma}{dt}|_E \approx 1.5 \cdot 10^{16} \text{s}^{-1}$ by curvature radiation with $\frac{d\gamma}{dt}|_{CR} \approx 8.5 \cdot 10^{-19} \gamma^4 \text{s}^{-1}$ one needs Lorentz factors of $\gamma \approx 3.6 \cdot 10^8$. Particles with smaller Lorentz factors experience only insignificant losses due to the γ^{-4} dependence of the curvature losses.

Inverse Compton scattering of relativistic particles on the thermal photons emitted by the hot neutron star surface (Sturmer 1995) limits the electrical acceleration of particles near the pulsar surface (Supper & Trümper, 2000). If one also includes the interaction of the charged particles with their own production of scattered photons one would expect an even more efficient braking to take place. Due to the curvature of their trajectories and we expect a fairly strong interaction of the particles with their own previously upscattered radiation within a few additional meters of travel, so that the effective energy density of the radiation field will increase quickly in the particle frame.

For an assumed complete dissipation of the available energy of the residual electric field near the surface we can estimate a effective temperature of

$$\begin{aligned} T &= \left(\frac{\epsilon_0 E_{\parallel}^2 (\xi - \frac{n}{n_{GJ}})^2}{a} \right)^{\frac{1}{4}} = \\ &= 7.3 \cdot 10^7 \text{K} \left(\xi - \frac{n}{n_{GJ}} \right)^{\frac{1}{2}} x^{-2} \end{aligned} \quad (13)$$

as a natural limit for the produced radiation. Here we use the blackbody constant $a = 7.566 \cdot 10^{-16} \text{Jm}^{-3} \text{K}^{-4}$ and $x = \frac{r}{r_{NS}}$ as the normalised distance from the surface. It is however unrealistic, to expect such efficient conversion of the available energy into radiation. Following Sturmer (1995) we expect dissipation scales for the dominant processes of the order of a few m in the vicinity of the neutron star surface. Again a simple estimate from a balance of the energy gain due to the electrical field on a dissipation scale λ will give us an improved temperature limit

$$\begin{aligned} T &= \left(e E_{\parallel} \left(1 - \frac{n}{n_{GJ}} \right) n \lambda \right)^{\frac{1}{4}} = \\ &= 4 \cdot 10^6 \text{K} x^{-1} \left(\left(1 - \frac{n}{n_{GJ}} \right) n \lambda \right)^{\frac{1}{4}} \end{aligned} \quad (14)$$

which turns out to be compatible with the observed x-ray temperatures of pulsars (Tsuruta 1998).

A more reliable description of the conditions near the pulsar surface can be attained by computing the

Lorentz factor of electrons with

$$\frac{d\gamma}{d\theta} = \frac{d\gamma}{d\theta}|_E + \frac{d\gamma}{d\theta}|_{CR} + \frac{d\gamma(T_{\text{eff}})}{d\theta}|_{\text{invCompt}} \quad (15)$$

together with the development of the energy density of the radiation field

$$\frac{dU}{d\theta} = -4U \frac{r_{\text{NS}}^2}{r(\theta, \theta_0)^3} - nm_e c^2 \frac{d\gamma(T_{\text{eff}})}{d\theta}|_{\text{invCompt}} \quad (16)$$

along a given field line parameterised by the colateral angles θ and θ_0 from the surface out to $10^3 r_{\text{NS}}$. The energy losses by the major processes outlined by Sturmer (1995) as there are: curvature radiation ($\frac{d\gamma}{d\theta}|_{CR}$), the Klein-Nishina correction as well as magnetic, resonant and non-resonant inverse Compton scattering, were included in the term $\frac{d\gamma(T_{\text{eff}})}{d\theta}|_{\text{invCompt}}$. The self-enhanced scattering from the radiation produced in lower levels of the magnetosphere was approximated by a local calculation of the effective temperature of the radiation bath $T_{\text{eff}} = (\frac{3U}{a})^{1/4}$. We did not calculate the exact spectrum of the radiation bath for reasons of efficiency. But we expect our crude approximation to be valid due to the curved trajectories of the electrons and the straight, tangential propagation of the emitted photons. After a short propagation, the scattering will not be back on, but at an slowly increasing angle. We repeated the calculations also without this enhancement effect, only to find, that the initial energy plateau did not extend quite as high up into the magnetosphere and the maximum energy achievable by the particles was a little higher. Because of the widely varying scales of the problem, the integration was performed for twelve segments of the trajectory using an Bulirsch-Stoer algorithm with decreasing resolution from one segment to the next (Press et al. 1992). The final values of each segment served as initial conditions for the subsequent ones. The results of each segment were carefully checked for their resolution independence, too coarse resolutions showing up as either oscillatory behaviour or non-convergence. Because of the enormous acceleration of a particle starting with $\gamma = 1$ at the surface, the first (surface-) segment had to be as short as 10 cm subdivided into 200 steps. Most interesting was the limiting case of $T = 2 \cdot 10^5$ K and $n = 0.9 n_{\text{GJ}}$. Fig. 7 provides a plot of the achieved Lorentz factors and effective temperatures as a function of distance from the surface. After travelling about 5 cm, the acceleration was balanced by non-resonant inverse Compton losses at $\gamma = 2.2 \cdot 10^5$. After that initial peaking of the

Lorentz factors, the particle energies decreased slowly down to $\gamma = 1.1 \cdot 10^5$ at $h = 20$ m accompanied by a slow rise of the effective temperature. That decrease is a consequence of self-enhancement and does not happen without it. Over the next 630 m the effective temperature rises quickly to 10^6 K and the Lorentz-factor reaches $1.3 \cdot 10^6$. Further, slower progress leads to a complete balance and saturation at $h = 1.3 \cdot 10^5$ m with $T = 2.9 \cdot 10^6$ K and $\gamma = 1.1 \cdot 10^8$. At $5.5 \cdot 10^4$ m curvature radiation takes over as the dominant energy loss mechanism. The Lorentz factors now decrease slowly down to $\gamma = 3.9 \cdot 10^7$ at a distance of $1.5 \cdot 10^7$ m.




Fig. 7.— The Lorentz factors (thin solid line: with self-enhanced inverse Compton scattering, thick solid: without enhanced scattering) and effective temperatures (dashed line) for enhanced scattering as a function of distance from the surface for $T = 2 \cdot 10^6$ K and $n = 0.9 \cdot n_{\text{GJ}}$.

A number of different calculations with varying initial parameters $n < 0.1 \cdot n_{\text{GJ}}$ and T were made, but showed no qualitative difference in the behaviour of their solutions. The case described can hence be taken as a reasonable representative of a "starved" system, having a low surface temperature and producing a strong source of X- and γ - rays with $T_{\text{eff}} \approx 3 \cdot 10^6$ K.

3.2. Pair production

Our calculations differ from those of Sturmer (1995) in as far as we did not *a priori* assume the existence of an inner polar gap. It is often unquestionably assumed that after about 100 m a particle creation avalanche creates a dense $n \gg n_{\text{GJ}}$ charge neutral plasma that shorts the available electric field. For an estimate of

the location of such an avalanche region we calculated the magnetic pair creation probability for curvature photons of a characteristic energy

$$E_{\text{curv}} = \frac{3ch\gamma^3}{4\pi R_c} \quad (17)$$

and Compton boosted thermal radiation having an energy of

$$E_{\text{comp}} = \gamma^2 k T_{\text{eff}} \quad (18)$$

The hard photons were assumed to be emitted tangentially and travelling outward from the field lines over a distance λ , their angle with the magnetic field lines $\Theta(\lambda, \theta, \theta_0)$ and the corresponding component of B perpendicular to the path of the photon

$$B_{\perp}(\lambda, \theta, \theta_0) = B_0 \cdot x^{-3} \sin \Theta(\lambda, \theta, \theta_0) \quad (19)$$

were explicitly calculated at each step along the trajectory of the accelerated charge. We used the local strength parameter

$$\chi(\gamma, \lambda, \theta, \theta_0) = \frac{\gamma}{2} \frac{B_{\perp}(\lambda, \theta, \theta_0)}{B_{\text{crit}}} \quad (20)$$

and Erber's (1966) Besselfunction approximation⁴ to calculate the local attenuation factor

$$\alpha_{\gamma B} = \frac{\alpha}{\lambda_e} \cdot \frac{0.16}{\chi} K_{1/3} \left(\frac{2}{3\chi} \right)^2 \quad (21)$$

with $\alpha = \frac{1}{137}$, the electron Compton wavelength $\lambda_e = \frac{\hbar}{m_e c}$ and the critical magnetic field $B_{\text{crit}} = \frac{m_e^2 c^2}{e\hbar} = 4.414 \cdot 10^9 \text{ T}$. $K_{1/3}$ stands for the modified Besselfunction of the third kind and order 1/3. Multiplying the attenuation factor $\alpha_{\gamma B}$ with the local steplength of the integration provides the local pair creation probability $p_{\gamma B}(\lambda, \theta, \theta_0)$ which in turn multiplied with the number of emitted photons, ie.

$$n_{\gamma} = \frac{2e^2}{9ch\varepsilon_0 R_c} \gamma = 2.1 \cdot 10^{-8} \gamma \text{ m}^{-1} \quad (22)$$

for curvature radiation, gives the local number of pairs created per accelerated charge⁵. We tested the

⁴Although that approximation involves more effort for its evaluation as compared to the commonly used exponential form, it accurate over a greater range of χ and, having no discontinuities, more useful for numerical work.

⁵Even under favourable conditions there will be only a small number of initial pairs produced. This is a consequence of the fact that $p_{\gamma B} \leq 1$ and that, at least for curvature radiation, the number of productive photons per unit length is very small. Because of the lower secondary and tertiary Lorentz factors, the conditions for efficient pair production in the form of daughter products tend to be even worse.

algorithm for the physically unrealistic case of missing inverse Compton braking and did indeed find a pair production zone at a height of 200 m above the pulsar surface.

However, of all our simulations that included non-resonant inverse Compton scattering, none showed $p_{\gamma B} > 0$ anywhere near the pulsar surface or within the simulated range of distances, neither for curvature photons nor for maximally boosted inverse Compton photons. This is due to the fact that the calculated Lorentz factors do not reach their highest values near the pulsar surface but further out, where the magnetic field has already fallen drastically due to its x^{-3} dependence.

4. Discussion and conclusions

We have seen that the surface temperature and the work function play a crucial role in determining the initial energy and density of the lower pulsar magnetosphere. Two scenarios are possible: Copious thermal emission of electrons leading to a "saturated" charged magnetosphere with $n = n_{\text{GJ}}$ and $E_{\parallel} = 0$. In that case we expect the inevitable potential drop and energy dissipation further away from the pulsar surface. Alternatively a "starved" magnetosphere, where $n < n_{\text{GJ}}$ is determined by the residual equilibrium field $(1 - \frac{n}{n_{\text{GJ}}})E_0$. Here our simulations differ from those made previously by others as we include the shielding of most of the $E_{\parallel}$ by the field emission produced particles together with the inverse Compton loss processes. By taking both physical effects into account, we find that conditions for pair creation cannot be met near the surface of ordinary rotating neutron stars that form most of the observed pulsars.

In the first case, we do not expect any Lorentz factors $\gamma > 1$. We have, however only described an unperturbed equilibrium atmosphere. The extensively studied fact that pulsar radio emission varies on timescales down to microseconds (Lange et al. (1998)) can be seen as an indication that such an atmosphere has inherent instabilities. Theoretical examples of instabilities in saturated magnetospheres were first given by Mestel et. al. (1985). Because of the strength of the compensated electric field, even smallest perturbations in the surface densities will be accompanied by (mildly) relativistic particles. Kunzl et. al. (1999) have described a scenario in which such perturbations lead to the radio emission via a drift induced free electron maser process. We expect that

most neutron stars to fall into that category, having surface temperatures higher than $2 \cdot 10^5 \text{K}$ combined with an $E_W \approx 400 \text{eV}$. If, as proposed by Pavlov & Zavlin (1998) neutron stars are surrounded by a gaseous atmosphere, we may expect easy liberation of charges in the form of electrons and ions, and a saturated magnetosphere is then unavoidable.

In the second category we would find cold neutron stars with a relativistic X-ray emitting lower magnetosphere, having effective temperatures of a few 10^6K . The achieved Lorentz factors are of the order of several 10^5 in the lower magnetosphere reaching a peak of $\gamma = 10^8$ at about $10r_{\text{NS}}$ and decrease monotonically to a few 10^7 at $1500r_{\text{NS}}$. A significant flow of X- and γ -rays can be expected in these circumstances, but the authors like to emphasise that radio emission models generally require much lower Lorentz factors ($\approx 10 - 1000$). The discussed loss processes are evidently incapable of braking the charges hard enough in order to achieve this, making difficult to explain any significant radio emission in these circumstances.

But the Lorentz factors in the region of strong magnetic fields are nevertheless too low to expect significant pair production and the subsequent formation of an inner gap in these cases.

But pulsars *are* radio and x-ray sources at the same time, this fact can only be reconciled with the arguments above by fully accepting the observational evidence that the magnetosphere is far from a homogeneous or steady state. A global investigation of instabilities in the current system of neutron star, magnetosphere and its outer boundary ought to be undertaken. Ideally, it would show that spatial or temporal variations in the magnetosphere would create the observed x-rays and the radio emission in different parts of the perturbation. Our calculations are meant to cover the possible extremes of the surface emission and the resultant charge densities and Lorentz-factors. Our terrestrial experience show us, that surface electron emission is affected by random variations of emission efficiency (formation of emitting islands) that are mainly responsible for the low-frequency part of a diode noise spectrum. It is not unthinkable that the thermal emission efficiency fluctuates over the polar cap. In that case we can expect local regions of the lower magnetosphere also to fluctuate individually between the two extreme cases described.

A few self critical remarks ought to be made: This study presents a drastically simplified scenario. We have omitted the effects of the pulsar's inclination

of the magnetic axis and possible surface inhomogeneities. Hence a greater spatial variation between the two extremes of starved and saturated charge liberation would be expected for real pulsars. We omitted a treatment that includes the effects of general relativity as i.e. was done by Muslimov and Harding (1997). If these effects are taken into account, fields and consequently Goldreich-Julian densities will be higher.

But due to the steep (exponential) characteristic of the surface emission processes, one expects a shift of the critical temperature by perhaps 10^5K . We also neglected resonances of the pair production process described by Daugherty & Harding, (1996) which lead to an enhancement of the pair production mechanism. The Lorentz factors compatible with inverse Compton braking however are by a factor of 10^3 lower than those constrained only by curvature radiation when pair production would be possible near the polar cap, and complete description of the problem would of course include both general relativistic effects and these resonances. A more detailed investigation of the physics of the upper surface layers would provide us with a more precise determination of E_W without the approximation of $E_W = E_F$. Furthermore, the simple approximation for T_{eff} does not incorporate the detailed spectral characteristics of the produced radiation, but the produced hard radiation will interact strongly with the accelerated electrons, we may even have underestimated the scattering losses.

We are lead to conclude, that the fact that we observe pulsars with X-ray temperatures of several 10^6K excludes the formation of an inner gap, and the fact that radio emission is also observed from such sources provides another argument against the high $n \gg n_{\text{GJ}}$ densities that are commonly expected from an inner avalanche region (Kunzl et al. 1998). We are also led to conclude, that observed pulsars are dominated by free charge emission with $n = n_{\text{GJ}}$, thus it might be advantageous to study the possible mechanisms for radio- and X-ray emission in that context.

REFERENCES

- Abrahams, A., & Shapiro, S., 1992, in *The Structure and Evolution of Neutron Stars*, Eds. D. Pines, R. Tamagaki & S. Tsuruta, Addison Wesley, p. 210
- Arendt, P.N., Eilek, J.A., 2000 in *Pulsar Astronomy – 2000 and beyond*, eds. Kramer, M., Wex, N., Wielebinski, R., IAU Colloq. **177**, p. 445

- Arons, J., Scharlemann, E., 1979, ApJ 231, 854
- Barnard, J.J., Arons, J., 1982, ApJ, 254, 713
- Daugherty, J.K. & Harding, A.K., 1996, ApJ, 458, 278
- Deutsch, A., 1955, Ann. Astrophys. 18, 1
- Dushman, S., 1930, Rev. Mod. Phys. 2, 470
- Edison, T.A., Engineering, 1884, 553
- Erber, T., 1966, Rev. Mod. Phys., 38, 626
- Flowers, E.G., Lee, J.F., Ruderman, M., Sutherland, G. Hillebrandt, W., & Müller, E., 1977, ApJ 215, 291
- Fowler, R.H., & Norheim, L.W., 1928, Roy. Soc. Proc. 119, 173
- Goldreich, P., & Julian, W.H., 1969, ApJ 157, 869
- Holloway, J.N., 1973, Nature, 246, 8-9
- Kijak, J. Gil, J., 1997, MNRAS, 288, 631
- Knoll, R. & Eichmeier, G., 1965, *Technische Elektronik*, Vol. 1, Springer Berlin
- Kramer, M., Xilouris, K.M., Jessner, A., Doroshenko, O., Wielebinski, R., 1997, ApJ, 488, 364
- Kramer, M., Xilouris, K.M., Jessner, A., von Hoensbroech, A., Lorimer, D., Wielebinski, R., Wolszczan, A., Camilo, F., 1998, ApJ, 501, 270
- Kunzl, T., Lesch, H., Jessner, A. & von Hoensbroech, A., 1998, ApJ, 505, L139
- Kunzl, T., Lesch, H. & Jessner, A., 1999, ApJ (submitted)
- Lange, Ch., Kramer, M., Wielebinski, R. & Jessner, A., 1998, A&A, 332, 111
- Lesch, H., Jessner, A., Kramer, M. & Kunzl, T., 1998 A&A, 332, L21
- Melrose, D.B., Gedalin, M.A., 1999, ApJ 521, 351
- Melrose, D.B., Gedalin, M.A., 2000, in Pulsar Astronomy – 2000 and beyond, eds. Kramer, M., Wex, N., Wielebinski, R., IAU Colloq. **177**, p. 381
- Mestel, L., Robertson, J.A., Wang, Y.-M. & Westfold, K.C., 1985, MNRAS, 217, 443
- Muslimov, A.G., & Tsygan, A.I., 1990, Sov. Astron. 34, 133
- Muslimov, A.G., & Tsygan, A.I., 1992, MNRAS 255, 61
- Muslimov, A.G., & Harding, A.K., 1997, ApJ 485, 735
- Pavlov, G.G. & Zavlin, V.E., 1998, in Neutron Stars and Pulsars, Ed. N. Shibasaki et al., Universal Academic Press, p. 327
- Press, W.H., Teukolsky, S.A., Vetterling, W.T. & Flannery, B.P., 1992, Numerical Recipes, Cambridge University Press
- Ruderman, M.A., Sutherland, P.G., 1975, ApJ 196, 51
- Schottky, W., 1914, Phys. Zeits. 15, 872
- Shibata, S., 1997, MNRAS, 287, 262
- Sturner, S.J., 1995, ApJ, 446, 292
- Supper, R., Trümper, J., 2000, A& A, 357, 301
- Usov, V.V., & Melrose, D.B., 1995, Aust. J. Phys., 48, 571
- Tsuruta, S., 1998, Phys. Rept., 292, 1
- Zhang, B. Harding, A.K., 2000, ApJ 532, 1150
- Zheleznyakov, V.V., 1996, Radiation Processes in Astrophysical Plasmas, Kluwer, Dordrecht